



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

Solution by the PROPOSER.

The only solution  $x$  divisible by  $p$  is seen to be  $\equiv 0 \pmod{p^n}$ . Let next  $x$  be prime to  $p$ . Then

$$x^{p^n-1} \equiv 1 \pmod{p^n}, \quad x^{p^{n-2}(p-1)} \equiv 1 \pmod{p^n},$$

the second from Fermat's theorem generalized, since  $\phi(p^n) = p^{n-1}(p-1)$ . Hence must  $x^{p-1} \equiv 1 \pmod{p^n}$ . To prove that the latter has precisely  $p-1$  distinct solutions  $\pmod{p^n}$ , we proceed by induction, the result being true for  $n=1$  by Fermat's theorem. Let therefore  $x_1, \dots, x_{p-1}$  be the distinct roots of  $x^{p-1} \equiv 1 \pmod{p^a}$ , whence  $x_i^p = x_i + r_i p^a$ , each  $r_i$  being a fixed integer. To determine the roots of  $x^{p-1} \equiv 1 \pmod{p^{a+1}}$ , we seek the integral values of  $m_i$  incongruent  $\pmod{p}$  such that  $x = x_i + m_i p^a$  satisfies  $x^p \equiv x \pmod{p^{a+1}}$ . But  $(x_i + m_i p^a)^p = x_i^p + \text{multiple of } p^{a+1}$ , by the Binomial Theorem. The condition is therefore

$$x_i + r_i p^a \equiv x_i + m_i p^a \pmod{p^{a+1}},$$

whence  $m_i \equiv r_i \pmod{p}$  and  $m_i$  is uniquely determined modulo  $p$ .

#### AVERAGE AND PROBABILITY.

148. Proposed by M. C. RORTY, Boston, Mass.

Assuming  $n$  points to fall at random upon a circle of circumference  $a$ , what is the probability of  $m$  or more points falling within a length  $b$  upon this circumference?

NOTE. This problem has practical application in determining the probability of accidental rushes of telephone calls as distinct from those rushes which are due to commercial causes. The solution for  $m$  or more points falling within a specified length  $b$  is known. The problem presented above differs from this in that a solution is required for *any* length  $b$ .

Solution by G. B. M. ZERR, A. M., Ph. D., Parsons, W. Va.

Let  $m+r$ , ( $r=0, 1, 2, 3, \dots, n-m$ ) be the number of points to fall on  $b$ .

$m+r$  points can be selected from  $n$  points in  $n! \div (m+r)!(n-m-r)!$  ways.

$$\therefore \text{Chance} = \frac{2n!}{(m+r)!(n-m-r)!} \cdot \frac{(b)^{m+r}(a-b)^{n-m-r}}{a^n}.$$

149. Proposed by L. C. WALKER, A.M., Professor of Mathematics, Colorado School of Mines, Golden, Col.

Three points are taken at random on the convex surface of a right cone. Find the probability that the section of the cone made by the plane passing through them is a complete ellipse.

Solution by G. B. M. ZERR, A. M., Ph. D., Parsons, W. Va.

Let  $ABC$  be a section of the cone,  $E$  one of the random points in the surface.  $AD=c$ ,  $GD=h$ ,  $BD=R$ ,  $EG=r$ . Through  $E$  pass the plane  $EC$ . Then the area of the surface, of which  $EBC$  is a projection

$$= \frac{\pi \sqrt{[h^2 + (R-r)^2]}}{R-r} [R^2 - (R+r)\sqrt{Rr}].$$